

EFFECTIVE DATE: 06|01|2023

POLICY LAST REVIEWED: 04|17|2024

OVERVIEW

Surgery for obesity, termed bariatric surgery, is a treatment for morbid obesity in patients who fail to lose weight with conservative measures. There are numerous different surgical techniques available. These different techniques have heterogenous mechanisms of action, with varying degrees of gastric restriction that creates a small gastric pouch, malabsorption of nutrients, and metabolic changes that result from gastric and intestinal surgery.

This policy only address services that are not covered for Medicare Advantage Plans and not medically necessary for Commercial Products.

For all other services, please refer to the Prior Authorization via Web-Based Tool for Procedures policy listed in the Related Policies section below.

MEDICAL CRITERIA

Not applicable

PRIOR AUTHORIZATION

Not applicable

POLICY STATEMENT

Medicare Advantage Plans

The bariatric surgery procedures listed below are considered not covered as the evidence is insufficient to determine the effects of the technology on health outcomes:

- Open adjustable gastric banding
- Open sleeve gastrectomy
- Open and laparoscopic vertical banding gastroplasty
- Gastric balloon for treatment of obesity
- Intestinal bypass

Note: Blue Cross & Blue Shield of Rhode Island (BCBSRI) must follow Centers for Medicare and Medicaid Services (CMS) guidelines, such as national coverage determinations or local coverage determinations for all Medicare Advantage Plans policies. Therefore, Medicare Advantage Plans policies may differ from Commercial products. In some instances, benefits for Medicare Advantage Plans may be greater than what is allowed by CMS.

Commercial Products

The bariatric surgery procedures listed below are considered not medically necessary for the treatment of morbid obesity in adults who have failed weight loss by conservative measures as the evidence is insufficient to determine the effects of the technology on health outcomes:

- Vertical-banded gastroplasty
- Gastric bypass using a Billroth II type of anastomosis (mini-gastric bypass)
- Biliopancreatic bypass without duodenal switch
- Long-limb gastric bypass procedure (i.e., >150 cm)
- Two-stage bariatric surgery procedures (e.g., sleeve gastrectomy as initial procedure followed by biliopancreatic diversion at a later time)

- Laparoscopic gastric plication
- Single anastomosis duodeno-ileal bypass with sleeve gastrectomy

The following endoscopic procedures are not medically necessary as a primary bariatric procedure or as a revision procedure (i.e., to treat weight gain after bariatric surgery to remedy large gastric stoma or large gastric pouches), as the evidence is insufficient to determine the effects of the technology on health outcomes:

- Insertion of the StomaphyX™ device
- Endoscopic gastroplasty
- Use of an endoscopically placed duodenojejunal sleeve
- Intra-gastric balloons
- Aspiration therapy device

COVERAGE

Benefits may vary between groups and contracts. Please refer to the appropriate Benefit Booklet, Evidence of Coverage or Subscriber Agreement for applicable surgery benefits/coverage or for applicable not medically necessary/not covered benefits/coverage.

BACKGROUND

Bariatric surgery is performed to treat obesity and obesity-related comorbid conditions. The first treatment of class III obesity is dietary and lifestyle changes. Although this strategy may be effective in some patients, only a few individuals with class III obesity can reduce and control weight through diet and exercise. Most patients find it difficult to comply with these lifestyle modifications on a long-term basis. When conservative measures fail, some patients may consider surgical approaches.

Intra-gastric balloon is unproven as a treatment for obesity. Further studies are needed to determine the safety and efficacy of intra-gastric balloon as a treatment option for obesity. Adverse effects associated with the intra-gastric balloon include gastric erosion, reflux, and obstruction (Fernandes, 2007). Currently, the available evidence in the published, peer-reviewed scientific literature is insufficient to establish the safety and efficacy of this procedure.

Vertical-banded gastroplasty was formerly one of the most common gastric restrictive procedures performed in the United States but has now been essentially replaced by other restrictive procedures due to high rates of revisions and reoperations. In this procedure, the stomach is segmented along its vertical axis. To create a durable reinforced and rate-limiting stoma at the distal end of the pouch, a plug of stomach is removed, and a propylene collar is placed through this hole and then stapled to itself. Because the normal flow of food is preserved, metabolic complications are uncommon. Complications include esophageal reflux, dilation, or obstruction of the stoma, with the latter two requiring reoperation. Dilation of the stoma is a common reason for weight regain. Vertical-banded gastroplasty may be performed using an open or laparoscopic approach.

Gastric Bypass Using a Billroth II Type of Anastomosis (mini-gastric bypass):

Recently, a variant of the gastric bypass, called the mini-gastric bypass, has been popularized. Using a laparoscopic approach, the stomach is segmented, similar to a traditional gastric bypass, but instead of creating a Roux-en-Y anastomosis, the jejunum is anastomosed directly to the stomach, similar to a Billroth II procedure. This unique aspect of this procedure is not based on its laparoscopic approach but rather the type of anastomosis used. It should also be noted that CPT code 43846 explicitly describes a Roux-en-Y gastroenterostomy, which is not used in the mini-gastric bypass.

Biliopancreatic Bypass (BPB) Without Duodenal Switch (CPT code 43847- gastric restrictive procedure, with gastric bypass for morbid obesity; with small intestine reconstruction to limit absorption):

Biliopancreatic bypass procedure (also known as the Scopinaro procedure) was developed and used extensively in Italy. It was designed to address some of the drawbacks of the original intestinal bypass procedures that have been abandoned due to unacceptable metabolic complications. Many of the complications were thought to be related to bacterial overgrowth and toxin production in the blind, bypassed segment. In contrast, BPB consists of a subtotal gastrectomy and diversion of the biliopancreatic juices into the distal ileum by a long Roux-en-Y procedure. The procedure consists of the following components:

- A distal gastrectomy induces a temporary early satiety and/or the dumping syndrome in the early postoperative period, both of which limit food intake.
- A 200-cm long “alimentary tract” consists of 200 cm of ileum connecting the stomach to a common distal segment.
- A 300- to 400-cm “biliary tract” connects the duodenum, jejunum, and remaining ileum to the common distal segment.
- A 50- to 100-cm “common tract” is where food from the alimentary tract mixes with biliopancreatic juices from the biliary tract. Food digestion and absorption, particularly of fats and starches, are therefore limited to this small segment of bowel, i.e., creating a selective malabsorption. The length of the common segment will influence the degree of malabsorption.

Because of the high incidence of cholelithiasis associated with the procedure, patients typically undergo an associated cholecystectomy.

Many potential metabolic complications are related to biliopancreatic bypass, including most prominently, iron deficiency anemia, protein malnutrition, hypocalcemia, and bone demineralization. Protein malnutrition may require treatment with total parenteral nutrition. In addition, there have been several case reports of liver failure resulting in death or liver transplant.

Single Anastomosis Duodenoileal Bypass with Sleeve Gastrectomy (SADI-S) (no specific CPT code):

No controlled trials of single anastomosis duodenoileal bypass with SG, (SADI-S), were identified. Some case series have been published that report on weight loss and other clinical outcomes up to 5 years post-surgery. One of the larger series was published in 2015 and reported on 97 patients with obesity and type 2 diabetes mellitus (DM). The authors reported that control of DM, defined as HgA1c <6.0%, was achieved in between 70% and 84% of patients at the different time points. Remission rates were higher for patients on oral therapy than those on insulin, and were higher in patients with a shorter duration of DM.

Long-Limb Gastric Bypass (le, >150 cm) (Gastric restrictive procedure with gastric bypass for morbid obesity; with small intestine reconstruction to limit absorption):

Recently, variations of gastric bypass procedures have been described, consisting primarily of long-limb Roux-en-Y procedures, which vary in the length of the alimentary and common limbs. For example, the stomach may be divided with a long segment of the jejunum (instead of ileum) anastomosed to the proximal gastric stump, creating the alimentary limb. The remaining pancreaticobiliary limb, consisting of stomach remnant, duodenum, and length of proximal jejunum, is then anastomosed to the ileum, creating a common limb of variable length in which the ingested food mixes with the pancreaticobiliary juices.

While the long alimentary limb permits absorption of most nutrients, the short common limb primarily limits absorption of fats. The stomach may be bypassed in a variety of ways, i.e., either by resection or stapling along the horizontal or vertical axis. Unlike the traditional gastric bypass, which is essentially a gastric restrictive procedure, these very long-limb Roux-en-Y gastric bypasses combine gastric restriction with some element of malabsorptive procedure, depending on the location of the anastomoses. Note that CPT code for gastric bypass (43846) explicitly describes a short limb (<150 cm Roux-en-Y gastroenterostomy, and thus would not apply to long-limb gastric bypass.

Two-Stage Procedure:

The evidence on the comparative efficacy of different bariatric surgery approaches consists largely of low-quality evidence, with a lack of long-term, high-quality randomized controlled trials (RCTs). Compared with gastric bypass, the evidence is sufficient to conclude that laparoscopic adjustable gastric banding is associated with lower short-term complications and lower medium- to long-term weight loss. The evidence is also

sufficient to conclude that sleeve gastrectomy has similar or lower short-term complications, with medium- to long-term weight loss that is somewhat less than for gastric bypass. The evidence on other types of bariatric surgery procedures is insufficient to form conclusions on the impact on health outcomes. For biliopancreatic bypass, the weight loss is similar or greater than gastric bypass but the complications rates, especially for nutritional complications, may also be higher. The evidence base for other types of procedures is insufficient to form conclusions.

Laparoscopic Gastric Plication (~~no specific CPT code~~):

Laparoscopic gastric plication is a bariatric surgery procedure that involves laparoscopic placement of sutures over the greater curvature (laparoscopic greater curvature plication) or anterior gastric region (laparoscopic anterior curvature plication) to create a tube-like stomach. The procedure involves two main steps, mobilization of the greater curvature of the stomach and suture plication of the stomach for achieving gastric restriction, but specifics of the technique are not standardized.

Endoscopic Procedures:

Endoscopic procedures (e.g., insertion of the StomaphyX™ device) as a primary bariatric procedure or as a revision procedure (i.e., to treat weight gain after bariatric surgery to remedy large gastric stoma or large gastric pouches): Some of these procedures use devices that are also being evaluated for endoscopic treatment of gastroesophageal reflux (GERD) (policy No. 2.01.38). The published data concerning use of these devices for treatment of regained weight is quite limited. Published case series have reported results using a number of different devices and procedures (including sclerosing injections) as treatment for this condition. The largest series found involved 28 patients treated with a sclerosing agent (sodium morrhuate). Reported trials that used one of the suturing devices had fewer than 10 patients. For example, Herron et al. reported on a feasibility study in animals. Thompson et al. reported on a pilot study with changes in anastomotic diameter and weight loss in 8 patients who had weight regain and dilated gastrojejunal anastomoses after RYGB. No comparative trials were identified; comparative trials are important because of the known association between an intervention and short-term weight loss. The StomaphyX™ device, which has been used in this approach, was cleared by the U.S. Food and Drug Administration (FDA) through the 510(k) process. It was determined to be equivalent to the EndoCinch™ system, which has 510(k) marketing clearance for endoscopic suturing for gastrointestinal tract surgery. In summary, the published scientific literature on use of these devices in patients who regain weight after bariatric surgery is very limited. No comparative studies were identified. These endoscopic procedures are considered investigational.

Bariatric Surgery in Patients with a BMI less than 35 KG/M2:

Limited evidence is available on bariatric surgery in patients with a BMI of less than 35 kg/m². Case series report a high rate of remission of diabetes in undergoing gastric bypass surgery, and this indication was judged to meet the TEC criteria in 2012. However, bariatric surgery for diabetes in patients with a BMI less than 35 is not currently considered standard of care and is not supported in current specialty society guidelines. For patients without diabetes, there is limited evidence on outcomes of surgery and no evidence that health outcomes are improved. As a result, bariatric surgery for patients with a BMI less than 35 is investigational.

The evidence on other types of bariatric surgery procedures is insufficient to form conclusions on the impact on health outcomes. For biliopancreatic bypass, the weight loss is similar or greater than gastric bypass but the complications rates, especially for nutritional complications, may also be higher. The evidence base for other types of procedures is insufficient to form conclusions.

For individuals who are preadolescent children with morbid obesity and receive bariatric surgery, the evidence includes no studies focused on this population. Relevant outcomes are overall survival, change in disease status, functional outcomes, health status measures, quality of life, and treatment related mortality and morbidity. Several studies of bariatric surgery in adolescents also included some children younger than 12 years old, but findings were not reported separately for preadolescent children. Moreover, clinical practice guidelines have recommended that bariatric surgery not be performed in preadolescent children. The evidence is insufficient to determine the effects of the technology on health outcomes.

CODING

Medicare Advantage Plans and Commercial Products

The following CPT code(s) are not covered for Medicare Advantage Plans and not medically necessary for Commercial Products:

- 43290** Esophagogastroduodenoscopy, flexible, transoral; with deployment of intragastric bariatric balloon
- 43842** Gastric restrictive procedure, without gastric bypass, for morbid obesity; vertical-banded gastroplasty
- 43843** Gastric restrictive procedure, without gastric bypass, for morbid obesity; other than vertical-banded gastroplasty
- 0813T** Esophagogastroduodenoscopy, flexible, transoral, with volume adjustment of intragastric bariatric balloon (New code 1/01/2024)

There are no specific CPT code(s) for the not covered/not medically necessary indications listed in this policy. Claims should be filed using the unlisted CPT code(s):

- 43659** Unlisted laparoscopy procedure, stomach
- 43999** Unlisted procedure, stomach

The following HCPCS code(s) is not separately reimbursed:

- S2083** Adjustment of gastric band diameter via subcutaneous port by injection or aspiration of saline

RELATED POLICIES

Non-Reimbursable Health Service Codes

Prior Authorization via Web-Based Tool for Procedures

Unlisted Procedures

PUBLISHED

Provider Update, June 2024

Provider Update, April 2023

Provider Update, May 2022

Provider Update, June 2021

Provider Update, September 2020

REFERENCES

1. Garvey WT, Mechanick JI, Brett EM, et al. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS AND AMERICAN COLLEGE OF ENDOCRINOLOGY COMPREHENSIVE CLINICAL PRACTICE GUIDELINES FOR MEDICAL CARE OF PATIENTS WITH OBESITYEXECUTIVE SUMMARYComplete Guidelines available at <https://www.aace.com/publications/guidelines>. Endocr Pract. Jul 2016; 22(7): 842-84. PMID 27472012
2. Centers for Disease Control and Prevention. Overweight & Obesity. Last Reviewed: June 3, 2022; <https://www.cdc.gov/obesity/basics/adult-defining.html>. Accessed March 29, 2024.
3. Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. JAMA. Oct 13 2004;292(14):1724-1737. PMID 15479938
4. Maggard MA, Shugarman LR, Suttorp M, et al. Meta-analysis: surgical treatment of obesity. Ann Intern Med. Apr 5 2005;142(7):547-559. PMID 15809466
5. Gomes-Rocha SR, Costa-Pinho AM, Pais-Neto CC, et al. Roux-en-Y Gastric Bypass Vs Sleeve Gastrectomy in Super Obesity: a Systematic Review and Meta-Analysis. Obes Surg. Jan 2022; 32(1): 170-185. PMID 34642872
6. Currie AC, Askari A, Fangueiro A, et al. Network Meta-Analysis of Metabolic Surgery Procedures for the Treatment of Obesity and Diabetes. Obes Surg. Oct 2021; 31(10): 4528-4541. PMID 34363144
7. Wilhelm SM, Young J, Kale-Pradhan PB. Effect of bariatric surgery on hypertension: a meta-analysis. Ann Pharmacother. Jun 2014;48(6):674-682. PMID 24662112

8. Ricci C, Gaeta M, Rausa E, et al. Early impact of bariatric surgery on type II diabetes, hypertension, and hyperlipidemia: a systematic review, meta-analysis and meta-regression on 6,587 patients. *Obes Surg.* Apr 2014;24(4):522-528. PMID 24214202
9. Cuspidi C, Rescaldani M, Tadic M, et al. Effects of bariatric surgery on cardiac structure and function: a systematic review and meta-analysis. *Am J Hypertens.* Feb 2014;27(2):146-156. PMID 24321879
10. Kwok CS, Pradhan A, Khan MA, et al. Bariatric surgery and its impact on cardiovascular disease and mortality: a systematic review and meta-analysis. *Int J Cardiol.* Apr 15 2014;173(1):20-28. PMID 24636546
11. Afshar S, Kelly SB, Seymour K, et al. The effects of bariatric surgery on colorectal cancer risk: systematic review and meta-analysis. *Obes Surg.* Oct 2014;24(10):1793-1799. PMID 25015708
12. Andersen JR, Aasprang A, Karlsten TI, et al. Health-related quality of life after bariatric surgery: a systematic review of prospective long-term studies. *Surg Obes Relat Dis.* Mar-Apr 2015; 11(2): 466-73. PMID 25820082
13. Arterburn DE, Olsen MK, Smith VA, et al. Association between bariatric surgery and long-term survival. *JAMA.* Jan 6 2015;313(1):62-70. PMID 25562267
14. Bower G, Toma T, Harling L, et al. Bariatric Surgery and Non-Alcoholic Fatty Liver Disease: a Systematic Review of Liver Biochemistry and Histology. *Obes Surg.* Dec 2015; 25(12): 2280-9. PMID 25917981
15. Cheung D, Switzer NJ, Ehmann D, et al. The impact of bariatric surgery on diabetic retinopathy: a systematic review and meta-analysis. *Obes Surg.* Sep 2015;25(9):1604-1609. PMID 25515499
16. Driscoll S, Gregory DM, Fardy JM, et al. Long-term health-related quality of life in bariatric surgery patients: A systematic review and meta-analysis. *Obesity (Silver Spring).* Jan 2016; 24(1): 60-70. PMID 26638116
17. Groen VA, van de Graaf VA, Scholtes VA, et al. Effects of bariatric surgery for knee complaints in (morbidly) obese adult patients: a systematic review. *Obes Rev.* Feb 2015;16(2):161-170. PMID 25487972
18. Hachem A, Brennan L. Quality of Life Outcomes of Bariatric Surgery: A Systematic Review. *Obes Surg.* Feb 2016; 26(2): 395-409. PMID 26494369
19. Lindekilde N, Gladstone BP, Lubeck M, et al. The impact of bariatric surgery on quality of life: a systematic review and meta-analysis. *Obes Rev.* Aug 2015; 16(8): 639-51. PMID 26094664
20. Lopes EC, Heineck I, Athaydes G, et al. Is Bariatric Surgery Effective in Reducing Comorbidities and Drug Costs? A Systematic Review and Meta-Analysis. *Obes Surg.* Sep 2015; 25(9): 1741-9. PMID 26112137
21. Ricci C, Gaeta M, Rausa E, et al. Long-term effects of bariatric surgery on type II diabetes, hypertension and hyperlipidemia: a meta-analysis and meta-regression study with 5-year follow-up. *Obes Surg.* Mar 2015;25(3):397-405. PMID 25240392
22. Yang XW, Li PZ, Zhu LY, et al. Effects of bariatric surgery on incidence of obesity-related cancers: a meta-analysis. *Med Sci Monit.* May 11 2015; 21: 1350-7. PMID 25961664
23. Madadi F, Jawad R, Mousati I, et al. Remission of Type 2 Diabetes and Sleeve Gastrectomy in Morbid Obesity: a Comparative Systematic Review and Meta-analysis. *Obes Surg.* Dec 2019; 29(12): 4066-4076. PMID 31655953
24. Yan G, Wang J, Zhang J, et al. Long-term outcomes of macrovascular diseases and metabolic indicators of bariatric surgery for severe obesity type 2 diabetes patients with a meta-analysis. *PLoS One.* 2019; 14(12): e0224828. PMID 31794559
25. Castellana M, Procino F, Biacchi E, et al. Roux-en-Y Gastric Bypass vs Sleeve Gastrectomy for Remission of Type 2 Diabetes. *J Clin Endocrinol Metab.* Mar 08 2021; 106(3): 922-933. PMID 33051679
26. Carmona MN, Santos-Sousa H, Lindeza L, et al. Comparative Effectiveness of Bariatric Surgeries in Patients with Type 2 Diabetes Mellitus and BMI ≥ 25 kg/m² : a Systematic Review and Network Meta-Analysis. *Obes Surg.* Dec 2021; 31(12): 5312-5321. PMID 34611827
27. Sjostrom L, Lindroos AK, Peltonen M, et al. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *N Engl J Med.* Dec 23 2004;351(26):2683-2693. PMID 15616203

28. Scopinaro N, Papadia F, Marinari G, et al. Long-term control of type 2 diabetes mellitus and the other major components of the metabolic syndrome after biliopancreatic diversion in patients with BMI < 35 kg/m². *Obes Surg.* Feb 2007;17(2):185-192. PMID 17476869
29. Sjostrom CD, Lissner L, Wedel H, et al. Reduction in incidence of diabetes, hypertension and lipid disturbances after intentional weight loss induced by bariatric surgery: the SOS Intervention Study. *Obes Res.* Sep 1999;7(5):477-484. PMID 10509605
30. Sjostrom L, Narbro K, Sjostrom CD, et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *N Engl J Med.* Aug 23 2007;357(8):741-752. PMID 17715408
31. Courcoulas AP, Christian NJ, Belle SH, et al. Weight change and health outcomes at 3 years after bariatric surgery among individuals with severe obesity. *JAMA.* Dec 11 2013;310(22):2416-2425. PMID 24189773
32. Arterburn D, Wellman R, Emiliano A, et al. Comparative Effectiveness and Safety of Bariatric Procedures for Weight Loss: A PCORnet Cohort Study. *Ann Intern Med.* Dec 04 2018; 169(11): 741-750. PMID 30383139
33. Arterburn DE, Johnson E, Coleman KJ, et al. Weight Outcomes of Sleeve Gastrectomy and Gastric Bypass Compared to Nonsurgical Treatment. *Ann Surg.* Dec 01 2021; 274(6): e1269-e1276. PMID 32187033
34. Wadden TA, Chao AM, Bahnsen JL, et al. End-of-Trial Health Outcomes in Look AHEAD Participants who Elected to have Bariatric Surgery. *Obesity (Silver Spring).* Apr 2019; 27(4): 581-590. PMID 30900413
35. Blue Cross Blue Shield Association Technology Evaluation Center (TEC). Laparoscopic adjustable gastric banding for morbid obesity. TEC Assessment. 2006;Vol 21:Tab 13.
36. Ibrahim AM, Thumma JR, Dimick JB. Reoperation and Medicare Expenditures After Laparoscopic Gastric Band Surgery. *JAMA Surg.* Sep 01 2017; 152(9): 835-842. PMID 28514487
37. Chakravarty PD, McLaughlin E, Whittaker D, et al. Comparison of laparoscopic adjustable gastric banding (LAGB) with other bariatric procedures; a systematic review of the randomised controlled trials. *Surgeon.* Jun 2012;10(3):172-182. PMID 22405735
38. Dixon JB, O'Brien PE, Playfair J, et al. Adjustable gastric banding and conventional therapy for type 2 diabetes: a randomized controlled trial. *JAMA.* Jan 23 2008;299(3):316-323. PMID 18212316
39. Gu L, Huang X, Li S, et al. A meta-analysis of the medium- and long-term effects of laparoscopic sleeve gastrectomy and laparoscopic Roux-en-Y gastric bypass. *BMC Surg.* Feb 12 2020; 20(1): 30. PMID 32050953
40. Han Y, Jia Y, Wang H, et al. Comparative analysis of weight loss and resolution of comorbidities between laparoscopic sleeve gastrectomy and Roux-en-Y gastric bypass: A systematic review and meta-analysis based on 18 studies. *Int J Surg.* Apr 2020; 76: 101-110. PMID 32151750
41. Sharples AJ, Mahawar K. Systematic Review and Meta-Analysis of Randomised Controlled Trials Comparing Long-Term Outcomes of Roux-En-Y Gastric Bypass and Sleeve Gastrectomy. *Obes Surg.* Feb 2020; 30(2): 664-672. PMID 31724116
42. Shenoy SS, Gilliam A, Mehanna A, et al. Laparoscopic Sleeve Gastrectomy Versus Laparoscopic Roux-en-Y Gastric Bypass in Elderly Bariatric Patients: Safety and Efficacy-a Systematic Review and Meta-analysis. *Obes Surg.* Nov 2020; 30(11): 4467-4473. PMID 32594469
43. Borgeraas H, Hofso D, Hertel JK, et al. Comparison of the effect of Roux-en-Y gastric bypass and sleeve gastrectomy on remission of type 2 diabetes: A systematic review and meta-analysis of randomized controlled trials. *Obes Rev.* Jun 2020; 21(6): e13011. PMID 32162437
44. Zhao H, Jiao L. Comparative analysis for the effect of Roux-en-Y gastric bypass vs sleeve gastrectomy in patients with morbid obesity: Evidence from 11 randomized clinical trials (meta-analysis). *Int J Surg.* Dec 2019; 72: 216-223. PMID 31756544
45. Lee Y, Doumouras AG, Yu J, et al. Laparoscopic Sleeve Gastrectomy Versus Laparoscopic Roux-en-Y Gastric Bypass: A Systematic Review and Meta-analysis of Weight Loss, Comorbidities, and Biochemical Outcomes From Randomized Controlled Trials. *Ann Surg.* Jan 01 2021; 273(1): 66-74. PMID 31693504
46. Xu C, Yan T, Liu H, et al. Comparative Safety and Effectiveness of Roux-en-Y Gastric Bypass and Sleeve Gastrectomy in Obese Elder Patients: a Systematic Review and Meta-analysis. *Obes Surg.* Sep 2020; 30(9): 3408-3416. PMID 32277330

47. Osland E, Yunus RM, Khan S, et al. Weight Loss Outcomes in Laparoscopic Vertical Sleeve Gastrectomy (LVSG) Versus Laparoscopic Roux-en-Y Gastric Bypass (LRYGB) Procedures: A Meta-Analysis and Systematic Review of Randomized Controlled Trials. *Surg Laparosc Endosc Percutan Tech.* Feb 2017; 27(1): 8-18. PMID 28145963
48. Osland EJ, Yunus RM, Khan S, et al. Five-Year Weight Loss Outcomes in Laparoscopic Vertical Sleeve Gastrectomy (LVSG) Versus Laparoscopic Roux-en-Y Gastric Bypass (LRYGB) Procedures: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Surg Laparosc Endosc Percutan Tech.* Dec 2020; 30(6): 542-553. PMID 32658120
49. Juodeikis Z, Brimas G. Long-term results after sleeve gastrectomy: A systematic review. *Surg Obes Relat Dis.* Apr 2017; 13(4): 693-699. PMID 27876332
50. Zhang Y, Wang J, Sun X, et al. Laparoscopic sleeve gastrectomy versus laparoscopic Roux-en-Y gastric bypass for morbid obesity and related comorbidities: a meta-analysis of 21 studies. *Obes Surg.* Jan 2015;25(1):19-26. PMID 25092167
51. Trastulli S, Desiderio J, Guarino S, et al. Laparoscopic sleeve gastrectomy compared with other bariatric surgical procedures: a systematic review of randomized trials. *Surg Obes Relat Dis.* Sep-Oct 2013;9(5):816-829. PMID 23993246
52. Brethauer SA, Hammel JP, Schauer PR. Systematic review of sleeve gastrectomy as staging and primary bariatric procedure. *Surg Obes Relat Dis.* Jul-Aug 2009;5(4):469-475. PMID 19632646
53. Hofso D, Fatima F, Borgeraas H, et al. Gastric bypass versus sleeve gastrectomy in patients with type 2 diabetes (Oseberg): a single-centre, triple-blind, randomised controlled trial. *Lancet Diabetes Endocrinol.* Dec 2019; 7(12): 912-924. PMID 31678062
54. Peterli R, Wölnerhanssen BK, Peters T, et al. Effect of Laparoscopic Sleeve Gastrectomy vs Laparoscopic Roux-en-Y Gastric Bypass on Weight Loss in Patients With Morbid Obesity: The SM-BOSS Randomized Clinical Trial. *JAMA.* Jan 16 2018; 319(3): 255-265. PMID 29340679
55. Salminen P, Helmiö M, Ovaska J, et al. Effect of Laparoscopic Sleeve Gastrectomy vs Laparoscopic Roux-en-Y Gastric Bypass on Weight Loss at 5 Years Among Patients With Morbid Obesity: The SLEEVEPASS Randomized Clinical Trial. *JAMA.* Jan 16 2018; 319(3): 241-254. PMID 29340676
56. Wölnerhanssen BK, Peterli R, Hurme S, et al. Laparoscopic Roux-en-Y gastric bypass versus laparoscopic sleeve gastrectomy: 5-year outcomes of merged data from two randomized clinical trials (SLEEVEPASS and SM-BOSS). *Br J Surg.* Jan 27 2021; 108(1): 49-57. PMID 33640917
57. Helmio M, Victorzon M, Ovaska J, et al. SLEEVEPASS: a randomized prospective multicenter study comparing laparoscopic sleeve gastrectomy and gastric bypass in the treatment of morbid obesity: preliminary results. *Surg Endosc.* Sep 2012;26(9):2521-2526. PMID 22476829
58. Karamanakos SN, Vagenas K, Kalfarentzos F, et al. Weight loss, appetite suppression, and changes in fasting and postprandial ghrelin and peptide-YY levels after Roux-en-Y gastric bypass and sleeve gastrectomy: a prospective, double blind study. *Ann Surg.* Mar 2008;247(3):401-407. PMID 18376181
59. Himpens J, Dapri G, Cadiere GB. A prospective randomized study between laparoscopic gastric banding and laparoscopic isolated sleeve gastrectomy: results after 1 and 3 years. *Obes Surg.* Nov 2006;16(11):1450-1456. PMID 17132410
60. Farrell TM, Haggerty SP, Overby DW, et al. Clinical application of laparoscopic bariatric surgery: an evidence- based review. *Surg Endosc.* May 2009;23(5):930-949. PMID 19125308
61. Skogar ML, Sundbom M. Duodenal Switch Is Superior to Gastric Bypass in Patients with Super Obesity when Evaluated with the Bariatric Analysis and Reporting Outcome System (BAROS). *Obes Surg.* Sep 2017; 27(9): 2308-2316. PMID 28439748
62. Strain GW, Gagner M, Inabnet WB, et al. Comparison of effects of gastric bypass and biliopancreatic diversion with duodenal switch on weight loss and body composition 1-2 years after surgery. *Surg Obes Relat Dis.* Jan- Feb 2007;3(1):31-36. PMID 17116424
63. Prachand VN, Davee RT, Alverdy JC. Duodenal switch provides superior weight loss in the super-obese (BMI > or =50 kg/m²) compared with gastric bypass. *Ann Surg.* Oct 2006;244(4):611-619. PMID 16998370
64. Strain GW, Torghabeh MH, Gagner M, et al. Nutrient Status 9 Years After Biliopancreatic Diversion with Duodenal Switch (BPD/DS): an Observational Study. *Obes Surg.* Jul 2017; 27(7): 1709-1718. PMID 28155056

65. Marceau P, Biron S, Hould FS, et al. Duodenal switch improved standard biliopancreatic diversion: a retrospective study. *Surg Obes Relat Dis.* Jan-Feb 2009;5(1):43-47. PMID 18440876
66. Cummings DE, Cohen RV. Bariatric/Metabolic Surgery to Treat Type 2 Diabetes in Patients With a BMI 35 kg/m². *Diabetes Care.* Jun 2016; 39(6): 924-33. PMID 27222550
67. Cummings DE, Rubino F. Metabolic surgery for the treatment of type 2 diabetes in obese individuals. *Diabetologia.* Feb 2018; 61(2): 257-264. PMID 29224190
68. Simonson DC, Vernon A, Foster K, et al. Adjustable gastric band surgery or medical management in patients with type 2 diabetes and obesity: three-year results of a randomized trial. *Surg Obes Relat Dis.* Dec 2019; 15(12): 2052-2059. PMID 31931977
69. Ikramuddin S, Billington CJ, Lee WJ, et al. Roux-en-Y gastric bypass for diabetes (the Diabetes Surgery Study): 2-year outcomes of a 5-year, randomised, controlled trial. *Lancet Diabetes Endocrinol.* Jun 2015; 3(6): 413-422. PMID 25979364
70. Liang Z, Wu Q, Chen B, et al. Effect of laparoscopic Roux-en-Y gastric bypass surgery on type 2 diabetes mellitus with hypertension: a randomized controlled trial. *Diabetes Res Clin Pract.* Jul 2013;101(1):50-56. PMID 23706413
71. Courcoulas AP, Belle SH, Neiberg RH, et al. Three-Year Outcomes of Bariatric Surgery vs Lifestyle Intervention for Type 2 Diabetes Mellitus Treatment: A Randomized Clinical Trial. *JAMA Surg.* Oct 2015; 150(10): 931-40. PMID 26132586
72. Courcoulas AP, Gallagher JW, Neiberg RH, et al. Bariatric Surgery vs Lifestyle Intervention for Diabetes Treatment: 5-Year Outcomes From a Randomized Trial. *J Clin Endocrinol Metab.* Mar 01 2020; 105(3): 866-76. PMID 31917447
73. Schauer PR, Bhatt DL, Kirwan JP, et al. Bariatric Surgery versus Intensive Medical Therapy for Diabetes - 5-Year Outcomes. *N Engl J Med.* Feb 16 2017; 376(7): 641-651. PMID 28199805
74. Mingrone G, Panunzi S, De Gaetano A, et al. Bariatric-metabolic surgery versus conventional medical treatment in obese patients with type 2 diabetes: 5 year follow-up of an open-label, single-centre, randomised controlled trial. *Lancet.* Sep 05 2015; 386(9997): 964-73. PMID 26369473
75. Wentworth JM, Playfair J, Laurie C, et al. Multidisciplinary diabetes care with and without bariatric surgery in overweight people: a randomised controlled trial. *Lancet Diabetes Endocrinol.* Jul 2014;2(7):545-552. PMID 24731535
76. Halperin F, Ding SA, Simonson DC, et al. Roux-en-Y gastric bypass surgery or lifestyle with intensive medical management in patients with type 2 diabetes: feasibility and 1-year results of a randomized clinical trial. *JAMA Surg.* Jul 2014;149(7):716-726. PMID 24899464
77. Scopinaro N, Adami GF, Papadia FS, et al. Effects of gastric bypass on type 2 diabetes in patients with BMI 30 to 35. *Obes Surg.* Jul 2014;24(7):1036-1043. PMID 24647849
78. Lanzarini E, Csendes A, Gutierrez L, et al. Type 2 diabetes mellitus in patients with mild obesity: preliminary results of surgical treatment. *Obes Surg.* Feb 2013;23(2):234-240. PMID 23054574
79. Boza C, Munoz R, Salinas J, et al. Safety and efficacy of Roux-en-Y gastric bypass to treat type 2 diabetes mellitus in non-severely obese patients. *Obes Surg.* Sep 2011;21(9):1330-1336. PMID 21744283
80. DePaula AL, Stival AR, DePaula CC, et al. Surgical treatment of type 2 diabetes in patients with BMI below 35: mid-term outcomes of the laparoscopic ileal interposition associated with a sleeve gastrectomy in 202 consecutive cases. *J Gastrointest Surg.* May 2012;16(5):967-976. PMID 22350720
81. Lee WJ, Wang W, Lee YC, et al. Effect of laparoscopic mini-gastric bypass for type 2 diabetes mellitus: comparison of BMI>35 and <35 kg/m². *J Gastrointest Surg.* May 2008;12(5):945-952. PMID 17940829
82. Matar R, Monzer N, Jaruvongvanich V, et al. Indications and Outcomes of Conversion of Sleeve Gastrectomy to Roux-en-Y Gastric Bypass: a Systematic Review and a Meta-analysis. *Obes Surg.* Sep 2021; 31(9): 3936-3946. PMID 34218416
83. Parmar CD, Gan J, Stier C, et al. One Anastomosis/Mini Gastric Bypass (OAGB-MGB) as revisional bariatric surgery after failed primary adjustable gastric band (LAGB) and sleeve gastrectomy (SG): A systematic review of 1075 patients. *Int J Surg.* Sep 2020; 81: 32-38. PMID 32738545
84. Brethauer SA, Kothari S, Sudan R, et al. Systematic review on reoperative bariatric surgery: American Society for Metabolic and Bariatric Surgery Revision Task Force. *Surg Obes Relat Dis.* Sep-Oct 2014;10(5):952-972. PMID 24776071

85. Dang JT, Vaughan T, Mocanu V, et al. Conversion of Sleeve Gastrectomy to Roux-en-Y Gastric Bypass: Indications, Prevalence, and Safety. *Obes Surg.* May 2023; 33(5): 1486-1493. PMID 36922465
86. Petrucciani N, Martini F, Benois M, et al. Revisional One Anastomosis Gastric Bypass with a 150-cm Biliopancreatic Limb After Failure of Adjustable Gastric Banding: Mid-Term Outcomes and Comparison Between One- and Two-Stage Approaches. *Obes Surg.* Dec 2021; 31(12): 5330-5341. PMID 34609712
87. Almalki OM, Lee WJ, Chen JC, et al. Revisional Gastric Bypass for Failed Restrictive Procedures: Comparison of Single-Anastomosis (Mini-) and Roux-en-Y Gastric Bypass. *Obes Surg.* Apr 2018; 28(4): 970-975. PMID 29101719
88. Sudan R, Nguyen NT, Hutter MM, et al. Morbidity, mortality, and weight loss outcomes after reoperative bariatric surgery in the USA. *J Gastrointest Surg.* Jan 2015;19(1):171-178; discussion 178-179. PMID 25186073
89. Catalano MF, Rudic G, Anderson AJ, et al. Weight gain after bariatric surgery as a result of a large gastric stoma: endotherapy with sodium morrhuate may prevent the need for surgical revision. *Gastrointest Endosc.* Aug 2007;66(2):240-245. PMID 17331511
90. Herron DM, Birkett DH, Thompson CC, et al. Gastric bypass pouch and stoma reduction using a transoral endoscopic anchor placement system: a feasibility study. *Surg Endosc.* Apr 2008;22(4):1093-1099. PMID 18027049
91. Thompson CC, Slattery J, Bundga ME, et al. Peroral endoscopic reduction of dilated gastrojejunal anastomosis after Roux-en-Y gastric bypass: a possible new option for patients with weight regain. *Surg Endosc.* Nov 2006;20(11):1744-1748. PMID 17024527
92. Eid GM, McCloskey CA, Eagleton JK, et al. StomaphyX vs a sham procedure for revisional surgery to reduce regained weight in Roux-en-Y gastric bypass patients: a randomized clinical trial. *JAMA Surg.* Apr 2014;149(4):372-379. PMID 24554030
93. Dakin GF, Eid G, Mikami D, et al. Endoluminal revision of gastric bypass for weight regain--a systematic review. *Surg Obes Relat Dis.* May-Jun 2013;9(3):335-342. PMID 23561960
94. Cohen RV, Oliveira da Costa MV, Charry L, et al. Endoscopic gastroplasty to treat medically uncontrolled obesity needs more quality data: A systematic review. *Surg Obes Relat Dis.* Jul 2019; 15(7): 1219-1224. PMID 31130406
95. Qi L, Guo Y, Liu CQ, et al. Effects of bariatric surgery on glycemic and lipid metabolism, surgical complication and quality of life in adolescents with obesity: a systematic review and meta-analysis. *Surg Obes Relat Dis.* Dec 2017; 13(12): 2037-2055. PMID 29079384
96. Black JA, White B, Viner RM, et al. Bariatric surgery for obese children and adolescents: a systematic review and meta-analysis. *Obes Rev.* Aug 2013;14(8):634-644. PMID 23577666
97. Treadwell JR, Sun F, Schoelles K. Systematic review and meta-analysis of bariatric surgery for pediatric obesity. *Ann Surg.* Nov 2008;248(5):763-776. PMID 18948803
98. Dumont PN, Blanchet MC, Gignoux B, et al. Medium- to Long-Term Outcomes of Gastric Banding in Adolescents: a Single-Center Study of 97 Consecutive Patients. *Obes Surg.* Jan 2018; 28(1): 285-289. PMID 29103071
99. Inge TH, Zeller MH, Jenkins TM, et al. Perioperative outcomes of adolescents undergoing bariatric surgery: the Teen-Longitudinal Assessment of Bariatric Surgery (Teen-LABS) study. *JAMA Pediatr.* Jan 2014;168(1):47-53. PMID 24189578
100. Olbers T, Beamish AJ, Gronowitz E, et al. Laparoscopic Roux-en-Y gastric bypass in adolescents with severe obesity (AMOS): a prospective, 5-year, Swedish nationwide study. *Lancet Diabetes Endocrinol.* Mar 2017; 5(3): 174-183. PMID 28065734
101. Willcox K, Brennan L. Biopsychosocial outcomes of laparoscopic adjustable gastric banding in adolescents: a systematic review of the literature. *Obes Surg.* Sep 2014;24(9):1510-1519. PMID 24849913
102. O'Brien PE, Sawyer SM, Laurie C, et al. Laparoscopic adjustable gastric banding in severely obese adolescents: a randomized trial. *JAMA.* Feb 10 2010;303(6):519-526. PMID 20145228
103. Nadler EP, Youn HA, Ren CJ, et al. An update on 73 US obese pediatric patients treated with laparoscopic adjustable gastric banding: comorbidity resolution and compliance data. *J Pediatr Surg.* Jan 2008;43(1):141-146. PMID 18206472

104. Manco M, Mosca A, De Peppo F, et al. The Benefit of Sleeve Gastrectomy in Obese Adolescents on Nonalcoholic Steatohepatitis and Hepatic Fibrosis. *J Pediatr.* Jan 2017; 180: 31-37.e2. PMID 27697327
105. Alqahtani AR, Elahmedi M, Abdurabu HY, et al. Ten-Year Outcomes of Children and Adolescents Who Underwent Sleeve Gastrectomy: Weight Loss, Comorbidity Resolution, Adverse Events, and Growth Velocity. *J Am Coll Surg.* Dec 2021; 233(6): 657-664. PMID 34563670
106. Greenstein RJ, Nissan A, Jaffin B. Esophageal anatomy and function in laparoscopic gastric restrictive bariatric surgery: implications for patient selection. *Obes Surg.* Apr 1998;8(2):199-206. PMID 9730394
107. Pilone V, Vitiello A, Hasani A, et al. Laparoscopic adjustable gastric banding outcomes in patients with gastroesophageal reflux disease or hiatal hernia. *Obes Surg.* Feb 2015;25(2):290-294. PMID 25030091
108. Kohn GP, Price RR, DeMeester SR, et al. Guidelines for the management of hiatal hernia. *Surg Endosc.* Dec 2013;27(12):4409-4428. PMID 24018762
109. Chen W, Feng J, Wang C, et al. Effect of Concomitant Laparoscopic Sleeve Gastrectomy and Hiatal Hernia Repair on Gastroesophageal Reflux Disease in Patients with Obesity: a Systematic Review and Meta-analysis. *Obes Surg.* Sep 2021; 31(9): 3905-3918. PMID 34254259
110. Mechanick JI, Apovian C, Brethauer S, et al. CLINICAL PRACTICE GUIDELINES FOR THE PERIOPERATIVE NUTRITION, METABOLIC, AND NONSURGICAL SUPPORT OF PATIENTS UNDERGOING BARIATRIC PROCEDURES - 2019 UPDATE: COSPONSORED BY AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS/AMERICAN COLLEGE OF ENDOCRINOLOGY, THE OBESITY SOCIETY, AMERICAN SOCIETY FOR METABOLIC BARIATRIC SURGERY, OBESITY MEDICINE ASSOCIATION, AND AMERICAN SOCIETY OF ANESTHESIOLOGISTS - EXECUTIVE SUMMARY. *Endocr Pract.* Dec 2019; 25(12): 1346-1359. PMID 31682518
111. Blonde L, Umpierrez GE, Reddy SS, et al. American Association of Clinical Endocrinology Clinical Practice Guideline: Developing a Diabetes Mellitus Comprehensive Care Plan-2022 Update. *Endocr Pract.* Oct 2022; 28(10): 923-1049. PMID 35963508
112. Department of Veterans Affairs/Department of Defense. Clinical Practice Guidelines. Management of Adult Overweight and Obesity (OBE) (2020). <https://www.healthquality.va.gov/guidelines/CD/obesity/>. Accessed April 2, 2024
113. Childerhose JE, Alsamawi A, Mehta T, et al. Adolescent bariatric surgery: a systematic review of recommendation documents. *Surg Obes Relat Dis.* Oct 2017; 13(10): 1768-1779. PMID 28958402
114. Armstrong SC, Bolling CF, Michalsky MP, et al. Pediatric Metabolic and Bariatric Surgery: Evidence, Barriers, and Best Practices. *Pediatrics.* Dec 2019; 144(6). PMID 31656225
115. Hampl SE, Hassink SG, Skinner AC, et al. Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents With Obesity. *Pediatrics.* Jan 09 2023. PMID 36622115
116. Michalsky M, Reichard K, Inge T, et al. ASMBS pediatric committee best practice guidelines. *Surg Obes Relat Dis.* Jan-Feb 2012;8(1):1-7. PMID 22030146
117. Pratt JSA, Browne A, Browne NT, et al. ASMBS pediatric metabolic and bariatric surgery guidelines, 2018. *Surg Obes Relat Dis.* Jul 2018; 14(7): 882-901. PMID 30077361
118. Eisenberg D, Shikora SA, Aarts E, et al. 2022 American Society for Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO): Indications for Metabolic and Bariatric Surgery. *Surg Obes Relat Dis.* Dec 2022; 18(12): 1345-1356. PMID 36280539
119. August GP, Caprio S, Fennoy I, et al. Prevention and treatment of pediatric obesity: an Endocrine Society clinical practice guideline based on expert opinion. *J Clin Endocrinol Metab.* Dec 2008;93(12):4576-4599. PMID 18782869
120. Styne DM, Arslanian SA, Connor EL, et al. Pediatric Obesity-Assessment, Treatment, and Prevention: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab.* Mar 01 2017; 102(3): 709-757. PMID 28359099
121. Centers for Medicare and Medicaid Services (CMS). Decision Memo for Bariatric Surgery for the Treatment of Morbid Obesity (CAG-00250R). 2006; <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=160>. Accessed April 1, 2024.
122. Gloy VL, Briel M, Bhatt DL, et al. Bariatric surgery versus non-surgical treatment for obesity: a systematic review and meta-analysis of randomised controlled trials. *BMJ.* Oct 22 2013;347:f5934. PMID 24149519

123. Puzifferri N, Roshek TB, 3rd, Mayo HG, et al. Long-term follow-up after bariatric surgery: a systematic review. *JAMA*. Sep 3 2014;312(9):934-942. PMID 25182102
124. Colquitt JL, Pickett K, Loveman E, et al. Surgery for weight loss in adults. *Cochrane Database Syst Rev*. 2014;8:CD003641. PMID 25105982
125. Kang JH, Le QA. Effectiveness of bariatric surgical procedures: A systematic review and network meta-analysis of randomized controlled trials. *Medicine (Baltimore)*. Nov 2017; 96(46): e8632. PMID 29145284
126. Park CH, Nam SJ, Choi HS, et al. Comparative Efficacy of Bariatric Surgery in the Treatment of Morbid Obesity and Diabetes Mellitus: a Systematic Review and Network Meta-Analysis. *Obes Surg*. Jul 2019; 29(7): 2180-2190. PMID 31037599
127. Cosentino C, Marchetti C, Monami M, et al. Efficacy and effects of bariatric surgery in the treatment of obesity: Network meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis*. Sep 22 2021; 31(10): 2815-2824. PMID 34348877

CLICK THE ENVELOPE ICON BELOW TO SUBMIT COMMENTS

This medical policy is made available to you for informational purposes only. It is not a guarantee of payment or a substitute for your medical judgment in the treatment of your patients. Benefits and eligibility are determined by the member's subscriber agreement or member certificate and/or the employer agreement, and those documents will supersede the provisions of this medical policy. For information on member-specific benefits, call the provider call center. If you provide services to a member which are determined to not be medically necessary (or in some cases medically necessary services which are non-covered benefits), you may not charge the member for the services unless you have informed the member and they have agreed in writing in advance to continue with the treatment at their own expense. Please refer to your participation agreement(s) for the applicable provisions. This policy is current at the time of publication; however, medical practices, technology, and knowledge are constantly changing. BCBSRI reserves the right to review and revise this policy for any reason and at any time, with or without notice. Blue Cross & Blue Shield of Rhode Island is an independent licensee of the Blue Cross and Blue Shield Association.

